



Primary Care COVID-19 Readiness Plan

The Western Public Health Unit have developed a 'COVID-19 Readiness Plan' to help GPs ensure patients at risk of developing severe disease are prepared and know how to link to appropriate testing and treatment in the event of COVID-19 infection. The tool can be completed with patients vulnerable to severe disease. It encourages patients to think about COVID-19 testing and treatment in advance, so that if it is required, they have a plan already in place.

Primary care practitioners can provide the completed tool to clinically vulnerable individuals to take home, giving them a tailored plan-of-action for rapid access to testing and treatment should symptoms develop.

The document can be downloaded <u>here</u> or found below.

Useful resources for primary care practitioners:

The Victorian COVID-19 therapies PBS prescriber helpline is available 7 days a week 8am to 5pm.

Phone: 03 8290 3801

The Victorian Department of Health has published the fact sheet: <u>COVID-19 mediations for at risk</u> <u>people who do not require oxygen</u>, which includes information about:

- Eligibility criteria for antiviral therapies.
- Clinical decision making.
- Prescribing considerations.
- Referral information for access to early therapies.







Your COVID-19 Readiness Plan

Date of		Clinia name
assessment:		Clinic name:
Name:		Clinic phone number:
Date of birth:		Usual GP:
		if you get COVID-19. Show this information to the right testing and treatment for you.
YOUR DOCTOR RECOMENDS If you get symptoms that cou support you need.		et as soon as you can. This will help you quickly get the care and
☐ Use a Rapid Antigen Test (I☐ Get a PCR test. The closest	•	
YOUR DOCTOR RECOMMEND If you have a positive test res antiviral medications as soon	ult on PCR or rapid antig	en test (RAT), your doctor recommends you get treatment with teps below.
Call your usual clinic on the place Tell them you have tested pospossible. Out of hours call:		our doctor has recommended you get treatment as soon as
Recommended antiviral	☐ Molnupiravir (Lagevrio TM) ☐ Nirmatrelvir+ritonavir (Paxlovid TM) ☐Other:	
Pharmacy where you can fill your prescription		
	ude severe shortness of	ALL TRIPLE ZERO (000) IMMEDIATELY TO GET HELP breath or difficulty breathing, chest pain, lips or face turning blue, usion or severe drowsiness.
Criteria for priority 1 (P1) PCI testing sites)	R testing (at State	Criteria for COVID-19 antiviral treatment eligibility
Unvaccinated or not up-to-date with vaccinations for COVID-19		 COVID-19 positive on rapid antigen or PCR test; AND Symptoms less than 5 days; AND

50 years and older with two additional risk factors Age 65 years or older Pregnant (including up to two weeks for developing severe disease; o 30 years and older AND identifying as Aboriginal or postpartum) Torres Strait Islander; Aboriginal and Torres Strait Islander 18 years and older with moderate to severe Underlying health conditions immuno compromisedClinician discretion OR 70 years and older; AND Symptoms less than 5 days OR asymptomatic and treatment is initiated as soon as possible after diagnosis is confirmed